

## Anti-AT2G20920 protein, N-terminal antibody

Catalog: PHY2209A

## **Product Information**

**Description:** Rabbit polyclonal antibody

Background: AT2G20920
Synonyms: AT2G20920

Immunogen: KLH-conjugated synthetic peptide (15 aa from N terminal section) derived from

Arabidopsis thaliana AT2G20920.

Form: Lyophilized

**Quantity**: 150 μg

Purification: Immunogen affinity purified

**Reconstitution:** Reconstitution with 150 µl of 0.01 M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

**Expected / apparent MW:** 30 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used


for immunization is 100% homologues with the sequence in *Brassica* rapa, *Brassica napus*, and 80-99% homologues with the sequence in *Vitis vinifera*, *Spinacia oleracea*, *Glycine max*, *Nicotiana tabacum*, *Medicago truncatula*, *Populus trichocarpa*, *Solanum tuberosum*, *Solanum lycopersicum*, *Cucumis sativus*, *Gossypium raimondii*, *Zea mays*, *Sorghum bicolor*, *Setaria viridis*, *Hordeum vulgare*, *Triticum* 



aestivum.

For more species homologues information, please contact tech support at <a href="tech@phytoab.com">tech@phytoab.com</a>.

## **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 43 kDa.

Electrophoresis: 12% SDS-PAGE

**Transfer:** blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or  $4^{\circ}$ C for 1 h.

**Primary antibody:** 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were

captured with CCD camera.