

## Anti-Photosystem I assembly protein Ycf3 antibody

Catalog: PHY5551S

## **Product Information**

Description:	Rabbit polyclonal antibody
Background:	A0A218N8W3
Synonyms:	YCF3
Immunogen:	KLH-conjugated synthetic peptide (18 aa from Central section) derived from
	Chlamydomonas reinhardtii YCF3 (A0A218N8W3).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Serum
	Peptide affinity form antibody available upon request at <u>info@phytoab.com</u> .
<b>Reconstitution:</b>	Reconstitution with 150 $\mu$ l of sterile water.
	"Note: please spin tube briefly prior to opening it to avoid any losses that might
	occur from lyophilized material adhering to the cap or sides of the tube".
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Storage:	12 months from date of receipt, -20 to -70 $^\circ\!\!\mathbb{C}$ as supplied.
	6 months, -20 to -70 $^\circ\!{ m C}$ under sterile conditions after reconstitution.
	1 month, 2 to 8 $^\circ\!\!\!\!^\circ$ under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4 $^\circ \! \mathbb{C}$ . Upon receipt, store it immediately at the
	temperature recommended above.

## **Application Information**

Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	20 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used
	for immunization is 80-99% homologues with the sequence in
	Gossypium raimondii, Brassica rapa, Populus trichocarpa, Zea mays,
	Brassica napus, Glycine max, Medicago truncatula, Solanum
	lycopersicum, Trit <mark>i</mark> cum aestivum, Cucumis sativus, Oryza sativa,
	Arabidopsis thaliana, Physcomitrium patens, Nicotiana tabacum,



Solanum tuberosum, Vitis vinifera, Spinacia oleracea, Hordeum vulgare, Panicum virgatum. For more species homologues information, please contact tech support at tech@phytoab.com.

## **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 41 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or  $4^{\circ}$ C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

**Secondary antibody:** 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured with CCD camera.

**Research Use Only**