

Anti-AT2G17972 antibody

Catalog: PHY2603A

Product Information

Description:	Rabbit polyclonal antibody
Background:	AT2G17972
Synonyms:	AT2G17972
Immunogen:	KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from
	Arabidopsis thaliana AT2G17972.
Form:	Lyophilized
Quantity:	150 µg
Purification:	Immunogen affinity purified
Reconstitution:	Reconstitution with 150 μ l of 0.01 M sterile PBS.
	"Note: please spin tube briefly prior to opening it to avoid any losses that might
	occur from lyophilized material adhering to the cap or sides of the tube".
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Storage:	12 months from date of receipt, -20 to -70 $^\circ C$ as supplied.
	6 months, -20 to -70 $^\circ\!\mathbb{C}$ under sterile conditions after reconstitution.
	1 month, 2 to 8 $^\circ C$ under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4 $^\circ\!{ m C}.$ Upon receipt, store it immediately at the
	temperature recommended above.

Application Information


Western Blot (1:1000-1:2000)
Note: Optimal dilutions/concentrations should be determined by the
end user.
18 / 13 kDa
Arabidopsis thaliana
Among species analyzed, the sequence of the synthetic peptide used
for immunization is 100% homologues with the sequence in <i>Panicum</i>
virgatum, Nicotiana tabacum, Oryza sativa, Sorghum bicolor, Setaria
viridis, Brassica napus, Brassica rapa, Solanum tuberosum, Solanum
<i>lycopersicum</i> , and 80-99% homologues with the sequence in
Gossypium raimondii, Zea mays, Populus trichocarpa, Hordeum

Research Use Only



vulgare, Triticum aestivum, Medicago truncatula, Glycine max, Vitis
vinifera, Cucumis sativus, Spinacia oleracea, Zea mays.
For more species homologues information, please contact tech
support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 43 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.

Research Use Only