

Anti-FLUCTUATING-LIGHT-ACCLIMATION PROTEIN1, C-terminal antibody

Catalog: PHY2540S

Product Information

Description: Rabbit polyclonal antibody

Background: Arabidopsis FLAP1 is associated with chloroplast thylakoid and envelope

membranes and that the *flap1* mutant shows delayed non-photochemical

quenching (NPQ) relaxation during induction of photosynthesis at moderate

light intensity.

Synonyms: FLAP1, FLUCTUATING-LIGHT-ACCLIMATION PROTEIN1

Immunogen: KLH-conjugated synthetic peptide (18 aa from C terminal section) derived from

Arabidopsis thaliana FLAP1 (AT1G54520).

Form: Lyophilized

Quantity:150 μgPurification:Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

Reconstitution: Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70° C as supplied.

6 months, -20 to -70 °C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:5000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 43 kDa


Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used



for immunization is 80-99% homologues with the sequence in Zea mays, Vitis vinifera, Glycine max, Oryza sativa, Triticum aestivum, Populus trichocarpa, Cucumis sativus, Gossypium raimondii, Nicotiana tabacum, Sorghum bicolor, Medicago truncatula, Solanum tuberosum, Hordeum vulgare, Setaria viridis, Brassica rapa, Solanum lycopersicum, Spinacia oleracea, Physcomitrium patens, Brassica napus.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 48 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP)

(Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.