

## Anti-Ferredoxin--NADP reductase, leaf isozyme 1, chloroplastic, C-terminal antibody

Catalog: PHY2233S

## **Product Information**

**Description:** Rabbit polyclonal antibody

**Background:** Two distinct ferredoxin-NADP(+)-oxidoreductase (FNR) isoforms were found in

chloroplasts of Arabidopsis thaliana, FNR-1 (AT5G66190) and FNR-2

(AT1G20020). The FNR proteins are present in both chloroplast stroma and

thylakoid membranes in chloroplasts but are more abundant in the stroma.

**Synonyms:** FNR1, ATLFNR1, FERREDOXIN-NADP(+)-OXIDOREDUCTASE 1, LEAF FNR

1, LEAF-TYPE CHLOROPLAST-TARGETED FNR 1, LFNR1

**Immunogen:** KLH-conjugated synthetic peptide (17 aa from C-terminal section) derived from

Arabidopsis thaliana FNR1 (AT5G66190).

Form: Lyophilized

**Quantity:** 150 μg **Purification:** Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

**Reconstitution:** Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70 °C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at  $4^{\circ}$ C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 40 / 34 kDa



Confirmed Reactivity: Arabidopsis thaliana

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used

for immunization is 100% homologues with the sequence in *Brassica* napus, *Brassica* rapa, and 80-99% homologues with the sequence in

Oryza sativa, Zea mays, Vitis vinifera, Sorghum bicolor, Gossypium

raimondii, Triticum aestivum, Panicum virgatum.

For more species homologues information, please contact tech

support at tech@phytoab.com.

## **Application Example**

Str: 12 µg stromal protein from Arabidopsis thaliana leaf.

Thy: Thylakoid membrane protein from Arabidopsis thaliana leaf containing 5 µg of chlorophyll.

Chl: 10 µl total chloroplast protein from Arabidopsis thaliana leaf.

TP: 18 µg total protein from Arabidopsis thaliana leaf.

Electrophoresis: 15% SDS-PAGE

**Transfer:** blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or 4°C for 1 h.

**Primary antibody:** 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG

H&L (HRP) (Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were

captured with CCD camera.

