

Anti-Ribonuclease J antibody

Catalog: PHY2322S

Product Information

Description: Rabbit polyclonal antibody

Background: Rnase J is a member of the metallo-beta-lactamase protein family that plays a

vital role in embryo morphogenesis and apical-basal pattern formation by

regulating chloroplast development.

Synonyms: Rnase J, ATRNJ, EMB2746, EMBRYO DEFECTIVE 2746, RIBONUCLEASE J,

RNJ

Immunogen: KLH-conjugated synthetic peptide (18 aa from C terminal section) derived from

Arabidopsis thaliana Rnase J (AT5G63420).

Form: Lyophilized

Quantity: 150 μg

Purification: Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

Reconstitution: Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70° C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8℃ under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 101 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used


for immunization is 100% homologues with the sequence in *Brassica* napus, *Brassica* rapa, and 80-99% homologues with the sequence in



Populus trichocarpa, Hordeum vulgare, Triticum aestivum, Cucumis sativus, Glycine max, Zea mays, Vitis vinifera, Medicago truncatula, Sorghum bicolor, Panicum virgatum, Gossypium raimondii, Oryza sativa, Setaria viridis.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 44 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4℃ for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP)

(Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with

CCD camera.