

Anti-Protein SCARECROW antibody

Catalog: PHY3959S

Product Information

Description: Rabbit polyclonal antibody

Background: SCR is a member of a novel family having similarity to DNA binding proteins

containing basic-leucine zipper regions; SCR appears to be a direct target of

SHR. SCR and SCR-LIKE 23 act redundantly in bundle sheath cell fate

specification.

Synonyms: SCR, SCARECROW, SGR1, SHOOT GRAVITROPISM 1

Immunogen: KLH-conjugated synthetic peptide (16 aa from C terminal section) derived from

Arabidopsis thaliana SCR (AT3G54220).

Form: Lyophilized

Quantity:150 μgPurification:Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

Reconstitution: Reconstitution with 150µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70° C as supplied.

6 months, -20 to -70 °C under sterile conditions after reconstitution.

1 month, 2 to 8℃ under sterile conditions after reconstitution.

Shipping: The product is shipped at 4° C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 71 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used


for immunization is 100% homologues with the sequence in Brassica

napus, Brassica rapa, Glycine max, Vitis vinifera.



For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 54 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4℃ for 1 h.

Primary antibody: 1:1000 dilution overnight at 4℃.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP)

(Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with

CCD camera.