

Anti-Meiotic recombination protein SPO11-1 antibody

Catalog: PHY1855S

Product Information

Description: Rabbit polyclonal antibody

Background: SPO11 is a key protein for promoting meiotic recombination, by generating

chromatin locus- and timing-specific DNA double-strand breaks (DSBs).

AtSPO11-1 is one of the three Arabidopsis homologues of the archaeal DNA

topoisomerase VIA subunit (topo VIA).

Synonyms: SPO11-1, ATSPO11-1, SPORULATION 11-1

Immunogen: KLH-conjugated synthetic peptide (18 aa from Central section) derived from

Arabidopsis thaliana SPO11-1 (AT3G13170).

Form: Lyophilized

Quantity:150 μgPurification:Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

Reconstitution: Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70° C as supplied.

6 months, -20 to -70 °C under sterile conditions after reconstitution.

1 month, 2 to 8℃ under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 42 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used


for immunization is 100% homologues with the sequence in Brassica



rapa, *Brassica napus*, and 80-99% homologues with the sequence in *Medicago truncatula*, *Hordeum vulgare*.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example:



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 38 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP)

(Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with

CCD camera.