

Anti-DEFECTIVE ACCUMULATION OF CYTOCHROME B6/F COMPLEX, C-terminal antibody

Catalog: PHY2793A

Product Information

Description:	Rabbit polyclonal antibody
Background:	DAC is a thylakoid membrane protein with two predicted transmembrane
	domains that is conserved from cyanobacteria to vascular plants. It is involved
	in the accumulation of the cytochrome b6/f complex.
Synonyms:	DAC, DEFECTIVE ACCUMULATION OF CYTOCHROME B6/F COMPLEX
Immunogen:	KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from
	Arabidopsis thaliana DAC (AT3G17930).
Form:	Lyophilized
Quantity:	150 μg
Purification:	Immunogen affinity purified
Reconstitution:	Reconstitution with 150 µl of 0.01M sterile PBS.
	"Note: please spin tube briefly prior to opening it to avoid any losses that might
	occur from lyophilized material adhering to the cap or sides of the tube".
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Storage:	12 months from date of receipt, -20 to -70 $^\circ C$ as supplied.
	6 months, -20 to -70 $^\circ\!\!\!\!\!^\circ$ under sterile conditions after reconstitution.
	1 month, 2 to 8 $^{\circ}$ C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4 $^\circ\!\!\!{}^\circ\!\!{}^\circ$. Upon receipt, store it immediately at the
	temperature recommended above.

Application Information


Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	21 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used for
	immunization is 100% homologues with the sequence in <i>Brassica napus,</i>
	<i>Brassica rap</i> a, and 80-99% homologues with the sequence in <i>Setaria</i>
	viridis, Triticum aestivum, Hordeum vulgare, Panicum virgatum, Zea

PhytoAB Inc.



mays, Oryza sativa, Sorghum bicolor, Cucumis sativus, Glycine max, Nicotiana tabacum, Solanum tuberosum, Solanum lycopersicum, Medicago truncatula, Gossypium raimondii.
For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example:



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 52 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4 °C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.

Research Use Only