

Anti-Ribosome associated endonuclease (CRB), C-terminal antibody

Catalog: PHY0158S

Product Information

Description:	Rabbit polyclonal antibody								
Background:	CSP41 protein has been shown to bind and cleave chloroplast RNA in vitro.								
	Arabidopsis thaliana genome encodes two copies of this protein CSP41a								
	(AT3G63140) and CSP41b (AT1G09340), and they interacts each other								
	physically. It has been suggested that CSP41a and CSP41b have a role in								
	chloroplast ribosomal RNA metabolism, most likely acting in the final steps of								
	23S rRNA maturation.								
Synonyms:	CSP41b, CHLOROPLAST RNA BINDING, CHLOROPLAST STEM-LOOP								
	BINDING PROTEIN OF 41 KDA, CRB, HETEROGLYCAN-INTERACTING								
	PROTEIN 1.3, HIP1.3								
Immunogen:	KLH-conjugated synthetic peptide (14 aa from C terminal section) derived from								
	Arabidopsis thaliana CSP41b (AT1G09340).								
Form:	Lyophilized								
Quantity:	150 µg								
Purification:	Serum								
	Peptide affinity form antibody available upon request at <u>info@phytoab.com</u> .								
Reconstitution:	Reconstitution with 150 μl of sterile water.								
	"Note: please spin tube briefly prior to opening it to avoid any losses that migh								
	occur from lyophilized material adhering to the cap or sides of the tube".								
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.								
Storage:	12 months from date of receipt, -20 to -70 $^\circ\!\!\!\mathrm{C}$ as supplied.								
	6 months, -20 to -70 $^\circ\mathrm{C}$ under sterile conditions after reconstitution.								
	1 month, 2 to 8 $^\circ\!\!\!\!^\circ$ under sterile conditions after reconstitution.								
Shipping:	The product is shipped at 4 $^\circ\!\!{ m C}$. Upon receipt, store it immediately at the								
	temperature recommended above.								

Application Information

Recommended Dilution:	Western Blo	t (1:10	000-1:4	4000)					
	Note: Optim	al dilu	itions/c	concent	rations	should	be det	ermined	by the

Research Use Only




Expected / apparent MW: Predicted Reactivity: end user.

43 kDa

Among species analyzed, the sequence of the synthetic peptide used for immunization is 100% homologues with the sequence in *Brassica napus*, *Brassica rapa*, *Panicum virgatum*, *Vitis vinifera*, *Solanum tuberosum*, *Solanum lycopersicum*, *Gossypium raimondii*, *Nicotiana tabacum*, *Zea mays*, and 80-99% homologues with the sequence in *Hordeum vulgare*, *Physcomitrium patens*, *Sorghum bicolor*, *Cucumis sativus*, *Populus trichocarpa*, *Triticum aestivum*, *Oryza sativa*, *Setaria viridis*, *Glycine max*, *Medicago truncatula*, *Spinacia oleracea*, *Chlamydomonas reinhardtii*.

For more species homologues information, please contact tech support at <u>tech@phytoab.com</u>.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 50 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.

Research Use Only