

Anti-NAD(P)H DEHYDROGENASE SUBUNIT 45 antibody

Catalog: PHY1772S

Product Information

Description: Rabbit polyclonal antibody

Background: NDH45 are novel nuclear-encoded subunits of the chloroplast NDH complex

and crucial both for the stable structure and function of the NDH complex.

NDH45 is involved in cyclic electron flow around photosystem I to produce ATP. The deficiency of either the NDH45 protein in respective mutant plants leads to

severe defects in both the accumulation and the function of the NDH complex.

Synonyms: NDH45, NAD(P)H DEHYDROGENASE SUBUNIT 45, NDF2,

NDH-DEPENDENT CYCLIC ELECTRON FLOW 1, PHOTOSYNTHETIC NDH

SUBCOMPLEX B 2, PNSB2

Immunogen: KLH-conjugated synthetic peptide (14 aa from C terminal section) derived from

Arabidopsis thaliana NDH45 (AT1G64770).

Form: Lyophilized

Quantity:150 μgPurification:Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

Reconstitution: Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70° as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8℃ under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 38 kDa

Research Use Only




Predicted Reactivity:

Among species analyzed, the sequence of the synthetic peptide used for immunization is 100% homologues with the sequence in *Brassica rapa*, *Brassica napus*, and 80-99% homologues with the sequence in *Glycine max*, *Vitis vinifera*, *Medicago truncatula*.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 43 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP)

(Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.