

Anti-bZIP transcription factor 11 antibody

Catalog: PHY1528A

Product Information

Description: Rabbit polyclonal antibody

Background: ZIP11 is a transcription factor that promotes POX1/PRODH1 (AT3G30775) in

response to hypoosmolarity stress. And it also regulates the ASN1

(AT3G47340) and POX2/PRODH2, which are involved in amino acid

metabolism.

Synonyms: BZIP11, ARABIDOPSIS THALIANA BASIC LEUCINE-ZIPPER 11, ATB2,

ATBZIP11, G-BOX BINDING FACTOR 6, GBF6

Immunogen: KLH-conjugated synthetic peptide (14 aa from central section) derived from

Arabidopsis thaliana BZIP11 (AT4G34590).

Form: Lyophilized

Quantity: 150 μg

Purification: Immunogen affinity purified

Reconstitution: Reconstitution with 150 µl of 0.01 M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70 °C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 18 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used


for immunization is 80-99% homologues with the sequence in Brassica rapa, Brassica napus, Medicago truncatula, Nicotiana



tabacum, Spinacia oleracea, Solanum tuberosum.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 43 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4℃ for 1 h.

Primary antibody: 1:1000 dilution overnight at 4℃.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP)

(Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with

CCD camera.