

Anti-3-dehydroquinate synthase antibody

Catalog: PHY2780A

Product Information

Description:	Rabbit polyclonal antibody	
Background:	AT3G28760	
Synonyms:	AT3G28760	
Immunogen:	KLH-conjugated synthetic peptide (16aa from central section) derived from	
	Arabidopsis thaliana AT3G28760.	
Form:	Lyophilized	
Quantity:	150 μg	
Purification:	Immunogen affinity purified	
Reconstitution:	Reconstitution with 150 μ l of 0.01 M sterile PBS.	
	"Note: please spin tube briefly prior to opening it to avoid any losses that might	
	occur from lyophilized material adhering to the cap or sides of the tube".	
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.	
Storage:	12 months from date of receipt, -20 to -70 $^\circ\mathrm{C}$ as supplied.	
	6 months, -20 to -70 $^\circ\!{ m C}$ under sterile conditions after reconstitution.	
	1 month, 2 to 8 $^\circ \!$	
Shipping:	The product is shipped at 4 $^\circ\!\mathrm{C}$. Upon receipt, store it immediately at the	
	temperature recommended above.	


Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	50 / 44 kDa
Confirmed Reactivity:	Arabidopsis thaliana
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used
	for immunization is 100% homologues with the sequence in <i>Brassica</i>
	rapa, Glycine max, Brassica napus, Cucumis sativus, Nicotiana
	tabacum, Solanum tuberosum, Oryza sativa, Solanum lycopersicum,
	and 80-99% homologues with the sequence in <i>Vitis vinifera</i> ,




Medicago truncatula, Zea mays, Hordeum vulgare, Sorghum bicolor, Setaria viridis, Panicum virgatum, Triticum aestivum, Populus trichocarpa, Spinacia oleracea, Chlamydomonas reinhardtii, Physcomitrium patens, Gossypium raimondii. For more species homologues information, please contact tech support at <u>tech@phytoab.com</u>.

Application Example Example1:



Example2:



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 54 kDa. **Electrophoresis:** 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4 °C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.

Research Use Only