

Anti-Eukaryotic translation initiation factor 3 subunit E antibody

Catalog: PHY7107S

Product Information

Description: Rabbit polyclonal antibody

Background: ORFX is a protein of unknown function. It has orthologous in other plants and

sequence is similar to E. coli ORFs orf154 and orf131, both of unknown function. So far, similar proteins are found only in plants and prokaryotes.

Synonyms: eIF3e, ATEIF3E-1, ATINT6, EIF3E, EUKARYOTIC TRANSLATION INITIATION

FACTOR 3E, INT-6, INT6, TIF3E1

Immunogen: KLH-conjugated synthetic peptide (18 aa from C terminal section) derived from

Arabidopsis thaliana eIF3e (AT3G57290).

Form: Lyophilized

Quantity:150 μgPurification:Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

Reconstitution: Reconstitution with 150 μl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Storage: Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

12 months from date of receipt, -20 to -70 °C as supplied.

6 months, -20 to -70 °C under sterile conditions after reconstitution.

1 month, 2 to 8℃ under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 52 kDa



Predicted Reactivity:

Among species analyzed, the sequence of the synthetic peptide used for immunization is 80-99% homologues with the sequence in Brassica napus, Brassica rapa, Glycine max, Medicago truncatula, Vitis vinifera, Glycine max, Gossypium raimondii, Spinacia oleracea, Nicotiana tabacum, Oryza sativa, Sorghum bicolor, Panicum virgatum, Setaria viridis, Populus trichocarpa, Cucumis sativus, Zea mays.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example

Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 43 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4° C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP)

(Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with

CCD camera.