

Anti-25.3 kDa vesicle transport protein, N-terminal antibody

Catalog: PHY0943S

Product Information

Description:	Rabbit polyclonal antibody
Background:	AtSec22 is a member of SEC22 Gene Family. AtSec22 is involved in vesicle
	trafficking from the ER to the Golgi complex.
Synonyms:	Sec22, ATSEC22, SEC221, SECRETION 22
Immunogen:	KLH-conjugated synthetic peptide (15 aa from N terminal section) derived from
	Arabidopsis thaliana Sec22 (AT1G11890).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Serum
	Peptide affinity form antibody available upon request at <u>info@phytoab.com</u> .
Reconstitution:	Reconstitution with 150 µl of ste <mark>r</mark> ile water.
	"Note: please spin tube briefly prior to opening it to avoid any losses that might
	occur from lyophilized material adhering to the cap or sides of the tube".
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Storage:	12 months from date of receipt, -20 to -70 $^\circ\!\!\mathbb{C}$ as supplied.
	6 months, -20 to -70 $^\circ\!\!\!\!\!^\circ$ under sterile conditions after reconstitution.
	1 month, 2 to 8 $^{\circ}$ C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4 $^\circ\!\mathrm{C}$. Upon receipt, store it immediately at the
	temperature recommended above.


Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	25 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used
	for immunization is 100% homologues with the sequence in <i>Brassica</i>
	<i>napus</i> , <i>Brassica rapa</i> , and 80-99% homologues with the sequence in
	Gossypium raimondii, Zea mays, Nicotiana tabacum, Oryza sativa



Indica Group, Triticum aestivum. For more species homologues information, please contact tech support at <u>tech@phytoab.com</u>.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 42 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.

Research Use Only