

Anti-Lysine-rich arabinogalactan protein 18 antibody

Catalog: PHY0965S

Product Information

Description: Rabbit polyclonal antibody

Background: ATAGP18 is a lysine-rich arabinogalactan-protein (AGP) and part of a

multi-gene family of glycoproteins with approx. 50 members. It falls into one subclass with AGP17 and AGP19, other lysine-rich AGPs. It is expressed in young leaves, shoots, roots and flowers and is active in the regulation of the

selection and survival of megaspores.

Synonyms: AGP18, ATAGP18, ARABINOGALACTAN PROTEIN 18

Immunogen: KLH-conjugated synthetic peptide (20 aa from C terminal section) derived from

Arabidopsis thaliana AGP18 (AT4G37450).

Form: Lyophilized

Quantity: 150 μg

Purification: Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

Reconstitution: Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70° C as supplied.

6 months, -20 to -70 °C under sterile conditions after reconstitution.

1 month, 2 to 8℃ under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 20 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used


for immunization is 80-99% homologues with the sequence in



Brassica rapa, Brassica napus.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide

for immunization and having a molecular mass of 54 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4℃ for 1 h.

Primary antibody: 1:1000 dilution overnight at 4° C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP)

(Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.