

Anti-30-kDa cleavage and polyadenylation specificity factor 30 antibody

Catalog: PHY3985S

Product Information

Description:	Rabbit polyclonal antibody
Background:	The CPSF proteins is the cleavage and polyadenylation specificity factor and is
	essential for the mRNA 3'-end processing. The Arabidopsis contains five CPSF
	proteins: CPSF160 (AT5G51660), CPSF100 (AT5G23880), CPSF73-I
	(AT1G61010), CPSF73-II (AT2G01730), and CPSF30 (AT1G30460).
Synonyms:	CPSF30, AtCPSF30, ARABIDOPSIS THALIANA CLEAVAGE AND
	POLYADENYLATION SPECIFICITY FACTOR 30, CLEAVAGE AND
	POLYADENYLATION SPECIFICITY FACTOR 30
Immunogen:	KLH-conjugated synthetic peptide (15 aa from Central section) derived from
	Arabidopsis thaliana CPSF30 (AT1G30460).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Serum
	Peptide affinity form antibody available upon request at <u>info@phytoab.com</u> .
Reconstitution:	Reconstitution with 150µl of sterile water.
	"Note: please spin tube briefly prior to opening it to avoid any losses that might
	occur from lyophilized material adhering to the cap or sides of the tube".
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Storage:	12 months from date of receipt, -20 to -70 $^\circ \!\! \mathbb{C}$ as supplied.
	6 months, -20 to -70 $^\circ\!\!\!\!\!^\circ$ under sterile conditions after reconstitution.
	1 month, 2 to 8 $^\circ C$ under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4 $^\circ\!\!{ m C}$. Upon receipt, store it immediately at the
	temperature recommended above.

Application Information


Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	70 kDa



Predicted Reactivity:

Among species analyzed, the sequence of the synthetic peptide used for immunization is 100% homologues with the sequence in *Solanum tuberosum*, *Cucumis sativus*, *Gossypium raimondii*, *Solanum lycopersicum*, *Nicotiana tabacum*, *Vitis vinifera*, *Brassica napus*, *Brassica rapa*, and 80-99% homologues with the sequence in *Glycine max*, *Spinacia oleracea*, *Oryza sativa*, *Medicago truncatula*, *Triticum aestivum*, *Hordeum vulgare*, *Zea mays*, *Sorghum bicolor*, *Panicum virgatum*, *Setaria viridis*, *Populus trichocarpa*, *Physcomitrium patens*. For more species homologues information, please contact tech support at <u>tech@phytoab.com</u>.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 50 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.