

## Anti-CBL-interacting protein kinase 9, C-terminal antibody

Catalog: PHY4454S

## **Product Information**

**Description:** Rabbit polyclonal antibody

Background: Os03g0126800

**Synonyms:** CIPK9

**Immunogen:** KLH-conjugated synthetic peptide (17 aa from C terminal section) derived from

Oryza sativa CIPK9 (Os03g0126800).

Form: Lyophilized

Quantity:150 μgPurification:Serum

Peptide affinity form antibody available upon request at <a href="mailto:info@phytoab.com">info@phytoab.com</a>.

**Reconstitution:** Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

**Storage:** 12 months from date of receipt, -20 to  $-70^{\circ}$ C as supplied.

6 months, -20 to -70 °C under sterile conditions after reconstitution.

1 month, 2 to 8<sup>°</sup>C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

**Expected / apparent MW:** 51 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used

for immunization is 100% homologues with the sequence in Glycine

max, Zea mays, Panicum virgatum, Setaria viridis, Gossypium


raimondii, Sorghum bicolor, Hordeum vulgare, Vitis vinifera, Cucumis



sativus, Medicago truncatula, and 80-99% homologues with the sequence in *Triticum aestivum*, *Populus trichocarpa*, *Solanum tuberosum*, *Nicotiana tabacum*, *Arabidopsis thaliana*, *Brassica rapa*, *Brassica napus*, *Solanum lycopersicum*.

For more species homologues information, please contact tech support at <a href="tech@phytoab.com">tech@phytoab.com</a>.

## **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide

for immunization and having a molecular mass of 50 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or  $4^{\circ}$ C for 1 h.

**Primary antibody:** 1:1000 dilution overnight at 4℃.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP)

(Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured with

CCD camera.