

## Anti-Vacuolar H+-ATPase subunit H antibody

Catalog: PHY7113S

## **Product Information**

**Description:** Rabbit polyclonal antibody

**Background:** V-ATPases are multisubunit membrane protein complexes that use the energy

provided by ATP hydrolysis to generate a proton gradient across various

intracellular and plasma membranes. Subunit H activates the ATPase activity of

the enzyme and couples ATPase activity to proton flow.

Synonyms: VHA-H

Immunogen: KLH-conjugated synthetic peptide (17 aa from N terminal section) derived from

Arabidopsis thaliana VHA-H (AT3G42050).

Form: Lyophilized

Quantity:150 μgPurification:Serum

Peptide affinity form antibody available upon request at <a href="mailto:info@phytoab.com">info@phytoab.com</a>.

**Reconstitution:** Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

**Storage:** 12 months from date of receipt, -20 to  $-70^{\circ}$ C as supplied.

6 months, -20 to -70 °C under sterile conditions after reconstitution.

1 month, 2 to 8℃ under sterile conditions after reconstitution.

**Shipping:** The product is shipped at  $4^{\circ}$ C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 50 kDa


Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used

for immunization is 80-99% homologues with the sequence in Brassica napus, Brassica rapa, Solanum tuberosum, Gossypium



raimondii, Nicotiana tabacum, Solanum lycopersicum, Populus trichocarpa, Vitis vinifera, Spinacia oleracea, Cucumis sativus. For more species homologues information, please contact tech support at <a href="tech@phytoab.com">tech@phytoab.com</a>.

## **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide

for immunization and having a molecular mass of 54 kDa.

**Electrophoresis:** 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or  $4^{\circ}$ C for 1 h.

**Primary antibody:** 1:1000 dilution overnight at  $4^{\circ}$ C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP)

(Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured with

CCD camera.