

Anti-Cytochrome b-c1 complex subunit Rieske-1/2, mitochondrial antibody

Catalog: PHY7121S

Product Information

Description: Rabbit polyclonal antibody

Background: Mitochondrial ubiquinol-cytochrome c oxidoreductase is also called complex III

and it accepts electron from quinol and transferred to cytochrome c. In

Arabidopsis mitochondria, ubiquinol-cytochrome c oxidoreductase contains 10 subunits. UCR1 (AT5G13430/AT5G13440) is one of the subunit of Complex III.

Synonyms: UCR1

Immunogen: KLH-conjugated synthetic peptide (14 aa from C terminal section) derived from

Arabidopsis thaliana UCR1-1 (AT5G13430) and UCR1-2 (AT5G13440).

Form: Lyophilized

Quantity: 150 μg

Purification: Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

Reconstitution: Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70° C as supplied.

6 months, -20 to -70 °C under sterile conditions after reconstitution.

1 month, 2 to 8℃ under sterile conditions after reconstitution.

Shipping: The product is shipped at 4° C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 30 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used


for immunization is 80-99% homologues with the sequence in Vitis



vinifera, Brassica rapa, Brassica napus, Hordeum vulgare, Triticum aestivum, Medicago truncatula.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 41 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4° C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP)

(Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.