Anti-ATP-dependent zinc metalloprotease FtsH 2, C-terminal antibody

Catalog: PHY5129A

Product Information

Description:	Rabbit polyclonal antibody
Background:	FtsH2 is acts as a processive, ATP-dependent zinc metallopeptidase for both cytoplasmic and membrane proteins. It plays a role in the quality control of integral membrane proteins.
Synonyms:	ftsH2
Immunogen:	KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from *Synechocystis sp*. PCC 6803 ftsH2 (slr0228).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Serum Peptide affinity form antibody available upon request at info@phytoab.com.
Reconstitution:	Reconstitution with 150 µl of sterile water. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70℃ as supplied. 6 months, -20 to -70℃ under sterile conditions after reconstitution. 1 month, 2 to 8℃ under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4℃. Upon receipt, store it immediately at the temperature recommended above.


Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the end user.
Expected / apparent MW:	68 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used for immunization is 80-99% homologues with the sequence in *Spinacia oleracea*, *Sorghum bicolor*, *Arabidopsis thaliana*, *Nicotiana*

tabacum, *Cucumis sativus*, *Panicum virgatum*, *Solanum lycopersicum*, *Vitis vinifera*, *Medicago truncatula*, *Populus trichocarpa*, *Brassica rapa*, *Brassica napus*, *Glycine max*, *Sorghum bicolor*, *Zea mays*, *Triticum aestivum*, *Hordeum vulgare*, *Setaria viridis*, *Vitis vinifera*, *Medicago truncatula*.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 45 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4℃ for 1 h.

Primary antibody: 1:1000 dilution overnight at 4℃.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.