

Anti-Probable protein phosphatase 2C 30, N-terminal antibody

Catalog: PHY0712A

Product Information

Description: Rabbit polyclonal antibody

Background: PP2C5 is a member of the PP2C family phosphatases. It acts as a MAPK

phosphatase that positively regulates seed germination, stomatal closure and ABA-inducible gene expression. The PP2C5 protein colocalizes and directly interacts with stress-induced MPK3 (AT3G45640), MPK4 (AT4G01370), and

MPK6 (AT2G43790), predominantly in the nucleus.

Synonyms: PP2C5, ATHPP2C5, PHOSPHATASE 2C5

Immunogen: KLH-conjugated synthetic peptide (15 aa from N terminal section) derived from

Arabidopsis thaliana PP2C5 (AT2G40180).

Form: Lyophilized

Quantity: 150 μg

Purification: Immunogen affinity purified

Reconstitution: Reconstitution with 150 µl of 0.01 M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70° C as supplied.

6 months, -20 to -70 °C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4° C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 43 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used


for immunization is 80-99% homologues with the sequence in



Brassica rapa, Brassica napus.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide

for immunization and having a molecular mass of 41 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4℃.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.