

# Anti-50S ribosomal protein L1p antibody

Catalog: PHY0421A

### **Product Information**

Description:	Rabbit polyclonal antibody	
Background:	50S ribosomal protein L1p (RP-L1p family) binds directly to 23S ribosomal	
	RNA.	
Synonyms:	RP-L1p, EMB3126, EMBRYO DEFECTIVE 3126, PLASTID RIBOSOMAL	
	PROTEIN L1, PRPL1	
Immunogen:	KLH-conjugated synthetic peptide (15 aa from N terminal section) derived from	
	Arabidopsis thaliana RP-L1p (AT3G63490).	
Form:	Lyophilized	
Quantity:	150 μg	
Purification:	Immunogen affinity purified	
Reconstitution:	Reconstitution with 150 μl of 0.01 M sterile PBS.	
	"Note: please spin tube briefly prior to opening it to avoid any losses that might	
	occur from lyophilized material adhering to the cap or sides of the tube".	
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.	
Storage:	12 months from date of receipt, -20 to -70 $^\circ\!\mathrm{C}$ as supplied.	
	6 months, -20 to -70 $^\circ \!$	
	1 month, 2 to 8 $^\circ\!\mathrm{C}$ under sterile conditions after reconstitution.	
Shipping:	The product is shipped at 4 $^\circ\!\mathrm{C}$ . Upon receipt, store it immediately at the	
	temperature recommended above.	

## **Application Information**


<b>Recommended Dilution:</b>	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	38 / 33 kDa
Confirmed Reactivity:	Arabidopsis thaliana
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used
	for immunization is 100% homologues with the sequence in <i>Brassica</i>
	napus, Vitis vinifera, Gossypium raimondii, Cucumis sativus,
	Medicago truncatula, Populus trichocarpa, Spinacia oleracea,



*Glycine max, Solanum tuberosum, Nicotiana tabacum, Brassica rapa, Solanum lycopersicum,* and 80-99% homologues with the sequence in *Physcomitrella patens, Oryza sativa, Triticum aestivum, Hordeum vulgare, Panicum virgatum, Sorghum bicolor, Zea mays, Setaria viridis.* 


For more species homologues information, please contact tech support at <u>tech@phytoab.com</u>.

## Application Example Example1:



Str: 15 μg stromal protein from *Arabidopsis thaliana*. **Electrophoresis:** 15% SDS- PAGE **Transfer:** blotting to NC (nitrocellulose) membrane for 1 h. **Blocking:** 5% skim milk at RT or 4°C for 1 h. **Primary antibody:** 1:1000 dilution overnight at 4°C. **Secondary antibody:** 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000) **Detection:** using chemiluminescence substrate and image were captured with CCD camera.

#### Example2:



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 45 kDa.
Electrophoresis: 12% SDS-PAGE
Transfer: blotting to NC (nitrocellulose) membrane for 1 h.
Blocking: 5% skim milk at RT or 4°C for 1 h.
Primary antibody: 1:1000 dilution overnight at 4°C.
Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).
Detection: using chemiluminescence substrate and image were captured

with CCD camera.

Research Use Only