

## Anti-Acetyl-coenzyme A carboxylase carboxyl transferase subunit alpha, chloroplastic antibody

Catalog: PHY3951S

## **Product Information**

**Description:** Rabbit polyclonal antibody

**Background:** CAC3 is the carboxyltransferase alpha subunit of acetyl-CoA carboxylase,

involved in de novo fatty acid biosynthesis.

**Synonyms:** CAC3, ACETYL CO-ENZYME A CARBOXYLASE CARBOXYLTRANSFERASE

**ALPHA SUBUNIT** 

**Immunogen:** KLH-conjugated synthetic peptide (15 aa from N terminal section) derived from

Arabidopsis thaliana CAC3 (AT2G38040).

Form: Lyophilized

Quantity:150 μgPurification:Serum

Peptide affinity form antibody available upon request at <a href="mailto:info@phytoab.com">info@phytoab.com</a>.

**Reconstitution:** Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Storage: Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

12 months from date of receipt, -20 to -70 °C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8℃ under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

**Recommended Dilution:** Western Blot(1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 85 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used


for immunization is 100% homologues with the sequence in *Brassica* napus, *Brassica* rapa, and 80-99% homologues with the sequence in



Medicago truncatula, Solanum tuberosum, Cucumis sativus, Glycine max, Spinacia oleracea, Solanum lycopersicum, Nicotiana tabacum, Populus trichocarpa, Gossypium raimondii, Vitis vinifera, Chlamydomonas reinhardtii, Physcomitrium patens.

For more species homologues information, please contact tech support at tech@phytoab.com.

## **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide

for immunization and having a molecular mass of 50 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4℃ for 1 h.

**Primary antibody:** 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.