

## Anti-Phycobilisome rod-core linker polypeptide CpcG antibody

Catalog: PHY5347S

## **Product Information**

Description:	Rabbit polyclonal antibody	
Background:	cpcG1	
Synonyms:	cpcG1, cpcG	
Immunogen:	KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from	
	Synechocystis sp. PCC 6803 cpcG1 (slr2051).	
Form:	Lyophilized	
Quantity:	150 µg	
Purification:	Serum	
	Peptide affinity form antibody available upon request at <u>info@phytoab.com</u> .	
Reconstitution:	onstitution: Reconstitution with 150 µl of sterile water.	
	"Note: please spin tube briefly prior to opening it to avoid any losses that might	
	occur from lyophilized material adhering to the cap or sides of the tube".	
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.	
Storage:	12 months from date of receipt, -20 to -70 $^\circ \!\! \mathbb C$ as supplied.	
	6 months, -20 to -70 $^\circ\!\!\!\!\!^\circ$ under sterile conditions after reconstitution.	
	1 month, 2 to 8 $^\circ C$ under sterile conditions after reconstitution.	
Shipping:	The product is shipped at 4 $^\circ\!\!{ m C}$ . Upon receipt, store it immediately at the	
	temperature recommended above.	


## **Application Information**

<b>Recommended Dilution:</b>	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	29 kDa
Predicted Reactivity:	For homologues with other species especially algae, please contact
	tech support at <u>tech@phytoab.com</u> .

Research Use Only



## **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 50 kDa. **Electrophoresis:** 12% SDS-PAGE **Transfer:** blotting to NC (nitrocellulose) membrane for 1 h. **Blocking:** 5% skim milk at RT or 4°C for 1 h. **Primary antibody:** 1:1000 dilution overnight at 4°C.

**Secondary antibody:** 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured with CCD camera.



**Research Use Only**