

Anti-Heat shock protein 90 antibody

Catalog: PHY0503

Product Information

Description:	Mouse monoclonal antibody
Background:	Hsp90 (heat shock protein 90) is a chaperone protein that assists other proteins
	to fold properly, stabilizes proteins against heat stress, and aids in protein
	degradation.
Synonyms:	HSP90, HSP81-2, Heat shock protein 90
Immunogen:	Recombinant protein HSP90 derived from Oryza sativa (Os09g0482100),
	which shows more than 90% identity with Arabidopsis thaliana HSP90 proteins
	(AT5G56030, AT5G56010, AT5G56000, AT5G52640).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Protein A purified
Reconstitution:	Reconstitution with 150 µl of 0.01 M sterile PBS.
	"Note: please spin tube bri <mark>efly prior to ope</mark> ning it to avoid any losses that might
	occur from lyophilized material adhering to the cap or sides of the tube".
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Storage:	12 months from date of receipt, -20 to -70 $^\circ \!\! \mathbb C$ as supplied.
	6 months, -20 to -70 $^\circ C$ under sterile conditions after reconstitution.
	1 month, 2 to 8° C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4 $^\circ\!\mathrm{C}$. Upon receipt, store it immediately at the
	temperature recommended above.

Application Information

Recommended Dilution:	Western Blot (1:1000-1:2500)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	80 kDa
Confirmed Reactivity:	Arabidopsis thaliana, Oryza sativa, Zea mays


Research Use Only



Predicted Reactivity:

For more species homologues information, please contact tech support at <u>tech@phytoab.com</u>.

Application Example



At-TP: 42 μg total protein from *Arabidopsis thaliana*. Os-TP: 25 μg, 100 μg total protein from *Oryza sativa*, respectively. Zm-TP: 25 μg, 100 μg total protein from *Zea mays*, respectively. **Electrophoresis:** 10% SDS-PAGE **Transfer:** blotting to NC (nitrocellulose) membrane for 1 h. **Blocking:** 5% skim milk at RT or 4°C for 1 h. **Primary antibody:** 1:5000 dilution overnight at 4°C. **Secondary antibody:** 1:5000 dilution using Goat Anti-Rabbit IgG H&L(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.

Research Use Only