

Anti-Uridylate kinase PUMPKIN, chloroplastic antibody

Catalog: PHY2842A

Product Information

Description: Rabbit polyclonal antibody

Background: PUMPKIN is a functional UMP Kinase located in the plastid that binds to group

Il intron plastid transcription products.

Synonyms: PUMPKIN, PLASTID 55 UMP KINASE

Immunogen: KLH-conjugated synthetic peptide (17 aa from central section) derived from

Arabidopsis thaliana PUMPKIN (AT3G18680).

Form: Lyophilized

Quantity: 150 μg

Purification: Immunogen affinity purified

Reconstitution: Reconstitution with 150 µl of 0.01M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70° C as supplied.

6 months, -20 to -70 °C under sterile conditions after reconstitution.

1 month, 2 to 8[°]C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 30 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used

for immunization is 100% homologues with the sequence in *Brassica napus*, *Brassica rapa*, *Spinacia oleracea*, and 80-99% homologues with the sequence in *Zea mays*, *Oryza sativa*, *Triticum aestivum*, *Hordeum vulgare*, *Sorghum bicolor*, *Panicum virgatum*, *Setaria viridis*, *Populus trichocarpa*, *Glycine max*, *Solanum lycopersicum*,


Research Use Only



Cucumis sativus, Gossypium raimondii, Nicotiana tabacum, Medicago truncatula, Solanum tuberosum, Vitis vinifera.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5ng and 10 ng recombinant protein containing the peptide for

immunization and having a molecular mass of 45 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4℃ for 1 h.

Primary antibody: 1:1000 dilution overnight at 4℃.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP)

(Cat# PHY6000)

Detection: using chemiluminescence substrate and image were captured with

CCD camera.