

Anti-Cytochrome f antibody

Catalog: PHY0488A

Product Information

Description: Rabbit polyclonal antibody

Background: Cytochrome f is a component of cytochrome b6/f complex, which mediates

electron transfer between photosystem II (PSII) and photosystem I (PSI), cyclic

electron flow around PSI, and state transitions.

Synonyms: PetA, PETA, PHOTOSYNTHETIC ELECTRON TRANSFER A

Immunogen: KLH-conjugated synthetic peptide (15 aa from central section) derived from

Arabidopsis thaliana PetA (ATCG00540).

Form: Lyophilized

Quantity: 150 μg

Purification: Immunogen affinity purified

Reconstitution: Reconstitution with 150 µl of 0.01 M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70 °C as supplied.

6 months, -20 to -70 °C under sterile conditions after reconstitution.

1 month, 2 to 8℃ under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 35 / 30 kDa

Confirmed Reactivity: Arabidopsis thaliana

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used

for immunization is 100% homologues with the sequence in *Brassica* napus, *Brassica* rapa, and 80-99% homologues with the sequence in

Zea mays, Oryza sativa, Nicotiana tabacum, Panicum virgatum,


Research Use Only



Sorghum bicolor, Gossypium raimondii, Glycine max, Medicago truncatula, Vitis vinifera, Solanum tuberosum, Cucumis sativus, Spinacia oleracea, Solanum lycopersicum, Hordeum vulgare, Populus trichocarpa, Triticum aestivum.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Thy-WT: thylakoid membrane protein from WT of *Arabidopsis thaliana* containing 0.25 μg and 0.5 μg of chlorophyll, respectively.

Thy-*Cyt b6f*: thylakoid membrane protein from Arabidopsis mutant with low accumulation of Cyt b6f complex containing 0.5 µg of chlorophyll.

Electrophoresis: 15% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4℃ for 1 h.

Primary antibody: 1:2000 dilution overnight at 4° C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG

H&L(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.