

## Anti-C-phycocyanin alpha subunit, C-terminal antibody

Catalog: PHY5242A

## **Product Information**

**Description:** Rabbit polyclonal antibody

**Background:** Phycocyanin alpha subunit, encoded by the *CpcA* (*sll1578*) in *Synechocystis* 

sp. PCC 6803.

Synonyms: CpcA

Immunogen: KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from

Synechocystis sp. PCC 6803 CpcA (sll1578).

Form: Lyophilized

**Quantity:** 150 μg

Purification: Immunogen affinity purified

**Reconstitution:** Reconstitution with 150 μl of 0.01 M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70 °C as supplied.

6 months, -20 to -70 °C under sterile conditions after reconstitution.

1 month, 2 to 8℃ under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 18 / 17 kDa


Confirmed Reactivity: Synechocystis sp. PCC 6803

Predicted Reactivity: For more species homologues information, please contact tech

support at tech@phytoab.com.



## **Application Example**



PCC 6803-TP: 31 µg and 62 µg total protein from Synechocystis sp. PCC 6803,

respectively.

Electrophoresis: 15% SDS-PAGE

**Transfer:** blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or 4°C for 1 h.

**Primary antibody:** 1:1000 dilution overnight at 4℃.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG

H&L(HRP) (Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured

with CCD camera.