

Anti-NAD(P)H-quinone oxidoreductase subunit 5, chloroplastic antibody

Catalog: PHY2979A

Product Information

Description: Rabbit polyclonal antibody

Background: NDHF is involved in the transfer of electrons from NAD(P)H: plastoquinone, via

FMN and iron-sulfur centers, to quinones in the photosynthetic chain and

possibly in a chloroplast respiratory chain.

Synonyms: NDHF

Immunogen: KLH-conjugated synthetic peptide (19 aa from N terminal section) derived from

Arabidopsis thaliana NDHF (ATCG01010).

Form: Lyophilized

Quantity: 150 μg

Purification: Immunogen affinity purified

Reconstitution: Reconstitution with 150 µl of 0.01 M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70° C as supplied.

6 months, -20 to -70 °C under sterile conditions after reconstitution.

1 month, 2 to 8℃ under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 85 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used

for immunization is 100% homologues with the sequence in


Medicago truncatula, Hordeum vulgare, Brassica rapa, Populus trichocarpa, Glycine max, Vitis vinifera, Cucumis sativus, Brassica



napus, Spinacia oleracea, Nicotiana tabacum, Leymus chinensis, Solanum tuberosum, Triticum aestivum, Zea mays, Sorghum bicolor, Setaria viridis, Solanum lycopersicum, Panicum virgatum, Gossypium raimondii, Oryza sativa, and 80-99% homologues with the sequence in *Physcomitrium patens*.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 15 ng and 25 ng recombinant proteincontaining the peptide for immunization and having a molecular mass of 40 kDa.

Electrophoresis: 12% SDS-PAGE.

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP)

(Cat# PHY6000).

Detection: using chemiluminescence substrate and image were

captured with CCD camera.