

Anti-CP47 subunit of Photosystem II antibody

Catalog: PHY0058A

Product Information

Description:	Rabbit polyclonal antibody	
Background:	CP47 is one of the components of the core complex of photosystem II (PSII). It	
	binds chlorophyll and helps catalyze the primary light-induced photochemical	
	processes of PSII. CP47 is encoded by ATCG00680.	
Synonyms:	PsbB	
Immunogen:	KLH-conjugated synthetic peptide (16 aa from C terminal section) derived from	
	Arabidopsis thaliana PsbB (ATCG00680).	
Form:	Lyophilized	
Quantity:	150 µg	
Purification:	Immunogen affinity purified	
Reconstitution:	Reconstitution with 150 μl of 0.01 M sterile PBS.	
	"Note: please spin tube briefly prior to opening it to avoid any losses that might	
	occur from lyophilized material adhering to the cap or sides of the tube".	
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.	
Storage:	12 months from date of receipt, -20 to -70 $^\circ\!\mathrm{C}$ as supplied.	
	6 months, -20 to -70 $^\circ \!\!\!\!\!^\circ \!\!\!^\circ$ under sterile conditions after reconstitution.	
	1 month, 2 to 8 $^\circ\!\mathrm{C}$ under sterile conditions after reconstitution.	
Shipping:	The product is shipped at 4 $^\circ\!\mathrm{C}$. Upon receipt, store it immediately at the	
	temperature recommended above.	

Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	56 kDa
Confirmed Reactivity:	Arabidopsis thaliana
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used
	for immunization is 100% homologues with the sequence in Solanum
	tuberosum, Physcomitrella patens, Spinacia oleracea, Medicago
	truncatula, Nicotiana tabacum, Zea mays, Populus trichocarpa, Vitis


Research Use Only



vinifera, Brassica rapa, Setaria viridis, Gossypium raimondii, Glycine max, Cucumis sativus, Brassica napus, Sorghum bicolor, and 80-99% homologues with the sequence in Oryza sativa, Hordeum vulgare, Triticum aestivum, Leymus chinensis, Chlamydomonas reinhardtii.

For more species homologues information, please contact tech support at <u>tech@phytoab.com</u>.

Application Example



Thy-WT: thylakoid membrane protein from WT of *Arabidopsis thaliana* containing 0.25 µg of chlorophyll. Thy-PS II mutant: thylakoid membrane protein from an Arabidopsis mutant with low accumulation of PS II containing 0.25 µg of chlorophyll. **Electrophoresis:** 15% SDS- PAGE **Transfer:** blotting to NC (nitrocellulose) membrane for 1 h. **Blocking:** 5% skim milk at RT or 4°C for 1 h. **Primary antibody:** 1:2000 dilution overnight at 4°C. **Secondary antibody:** 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000) **Detection:** using chemiluminescence substrate and image were captured with CCD camera.

Research Use Only