

# Anti-AT1G62780, C-terminal antibody

Catalog: PHY2622S

### **Product Information**

**Description:** Rabbit polyclonal antibody

Background: AT1G62780

Synonyms: AT1G62780

Immunogen: KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from

Arabidopsis thaliana (AT1G62780).

Form: Lyophilized

Quantity:150 μgPurification:Serum

Peptide affinity form antibody available upon request at <a href="mailto:info@phytoab.com">info@phytoab.com</a>.

**Reconstitution:** Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

**Recommended Dilution:** Western Blot (1:1000-1:5000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 27 / 26 kDa

Confirmed Reactivity: Arabidopsis thaliana

**Predicted Reactivity:** Among species analyzed, the sequence of the synthetic peptide used

for immunization is 80-99% homologues with the sequence in Brassica

rapa, Brassica napus, Gossypium raimondii, Glycine max, Vitis

vinifera, Physcomitrium patens, Cucumis sativus.



For more species homologues information, please contact tech support at <a href="tech@phytoab.com">tech@phytoab.com</a>.

### **Application Example**

#### **Example 1**

Chl: 4 µl total chloroplast protein from Arabidopsis thaliana.

Electrophoresis: 15% SDS-PAGE

**Transfer:** blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or  $4^{\circ}$ C for 1 h.

**Primary antibody:** 1:5000 dilution overnight at 4°C.


Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000)

Detection: using chemiluminescence substrate and image were captured with CCD

camera.

### Example 2



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide

for immunization and having a molecular mass of 47 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

**Primary antibody:** 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.