

Anti-Os02g0600200 protein, C-terminal antibody

Catalog: PHY4349S

Product Information

Description: Rabbit polyclonal antibody

Background: Os02g0600200

Synonyms: Os02g0600200

Immunogen: KLH-conjugated synthetic peptide (16 aa from C terminal section) derived from

Oryza sativa Os02g0600200.

Form: Lyophilized

Quantity:150 μgPurification:Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

Reconstitution: Reconstitution with 150 μl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70°C as supplied.

6 months. -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:5000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 30 / 24 kDa

Confirmed Reactivity: Oryza sativa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used


for immunization is 80-99% homologues with the sequence in Sorghum bicolor, Zea mays, Panicum virgatum, Setaria viridis,

Triticum aestivum, Hordeum vulgare.



For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example Example1:



Os-Chl: 5 µl total chloroplast protein from Oryza sativa.

Electrophoresis: 15% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:5000 dilution overnight at 4°C.


Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG

H&L(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were

captured with CCD camera.

Example2:



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide

for immunization and having a molecular mass of 46 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4℃.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.