

Anti-Photosynthetic NDH subunit of subcomplex B 4, chloroplastic antibody

Catalog: PHY7270S

Product Information

Description: Rabbit polyclonal antibody

Background: PNSB4 is likely a subunit of the chloroplast NAD(P)H dehydrogenase complex,

it is involved in PSI cyclic electron transport. It located on the thylakoid

membrane.

Synonyms: PNSB4, NDF6, NDH DEPENDENT FLOW 6, PHOTOSYNTHETIC NDH

SUBCOMPLEX B 4

Immunogen: KLH-conjugated synthetic peptide (16 aa from C terminal section) derived from

Arabidopsis thaliana PNSB4 (AT1G18730).

Form: Lyophilized

Quantity:150 μgPurification:Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

Reconstitution: Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70° C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 20 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide


used for immunization is 100% homologues with the sequence in



Brassica rapa, Brassica napus, and 80-99% homologues with the sequence in Spinacia oleracea, Nicotiana tabacum, Solanum tuberosum, Solanum lycopersicum, Medicago truncatula, Vitis vinifera.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 45 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG

H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were

captured with CCD camera.