

Anti-YCF54 antibody

Product Information

Description:	Rabbit polyclonal antibody	
Background:	YCF54 is located in chloroplast, it involved in biological process.	
Synonyms:	YCF54, CGL78, EMB3143, EMBRYO DEFECTIVE 3143	
Immunogen:	KLH-conjugated synthetic peptide (17 aa from Central section) derived from	
	Arabidopsis thaliana YCF54 (AT5G58250).	
Form:	Lyophilized	
Quantity:	150 µg	
Purification:	Serum	
Reconstitution:	Reconstitution with 150 µl of sterile water.	
	"Note: please spin tube briefly prior to opening it to avoid any losses that might	
	occur from lyophilized material adhering to the cap or sides of the tube".	
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.	
Storage:	12 months from date of receipt, -20 to -70 °C as supplied.	
	6 months, -20 to -70 $^\circ\!{ m C}$ under sterile conditions after reconstitution.	
	1 month, 2 to 8 $^\circ\!\!\!\!^\circ$ under sterile conditions after reconstitution.	
Shipping:	The product is shipped at 4 $^\circ\!\!\mathbb{C}.$ Upon receipt, store it immediately at the	
	temperature recommended above.	

Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected/apparent MW:	24 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used
	for immunization is 100% homologues with the sequence in <i>Brassica</i>
	napus, Brassica rapa, Glycine max, Nicotiana tabacum, and 80-99%
	homologues with the sequence in Solanum tuberosum, Medicago
	truncatula, Solanum lycopersicum, Sorghum bicolor, Setaria viridis,
	Panicum virgatum, Zea mays, Oryza sativa Japonica Group, Triticum
	aestivum, Hordeum vulgare, Physcomitrium patens, Vitis vinifera,

Research Use Only




Gossypium raimondii.

captured with CCD camera.

For more species homologues information, please contact tech support at <u>tech@phytoab.com</u>.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 47 kDa. **Electrophoresis:** 12% SDS-PAGE **Transfer:** blotting to NC (nitrocellulose) membrane for 1 h. **Blocking:** 5% skim milk at RT or 4°C for 1 h. **Primary antibody:** 1:1000 dilution overnight at 4°C. **Secondary antibody:** 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000). **Detection:** using chemiluminescence substrate and image were

PHY0663S

Research Use Only