

Anti-AT5G14105 antibody

Catalog: PHY0552S

Product Information

Description: Rabbit polyclonal antibody

Background: AT5G14105

Synonyms: AT5G14105

Immunogen: KLH-conjugated synthetic peptide (14 aa from Central section) derived from

Arabidopsis thaliana AT5G14105.

Form: Lyophilized

Quantity: $150 \mu g$ Purification: Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

Reconstitution: Reconstitution with 150 μl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70° C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4 °C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 8 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide

used for immunization is 100% homologues with the sequence in

Populus trichocarpa, and 80-99% homologues with the sequence in Brassica napus, Brassica rapa, Solanum lycopersicum, Glycine max,


Gossypium raimondii, Medicago truncatula, Cucumis sativus,



Nicotiana tabacum, Solanum tuberosum, Zea mays, Panicum virgatum, Setaria viridis, Oryza sativa, Vitis vinifera, Sorghum bicolor, Spinacia oleracea.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 50 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4 °C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.