

## Anti-Translocase of chloroplast 33, chloroplastic antibody

Catalog: PHY3495A

## **Product Information**

Description:	Rabbit polyclonal antibody
Background:	TOC33 (AT1G02280) is a component of the chloroplast protein import
	machinery, which consists of three core components, namely the pore-forming
	Toc75 (AT1G35860, AT3G46740, AT4G09080, AT5G19620) and two GTPases
	Toc33 (AT1G02280) and Toc159 (AT4G02510). Toc33 and Toc159 may
	regulate precursor protein perception and their transport to the translocation
	pore.
Synonyms:	TOC33, ATTOC33, PLASTID PROTEIN IMPORT 1, PPI1, TRANSLOCON AT
	THE OUTER ENVELOPE MEMBRANE OF CHLOROPLASTS 33
Immunogen:	KLH-conjugated synthetic peptide (14 aa from C terminal section) derived from
	Arabidopsis thaliana TOC33 (AT1G02280).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Immunogen affinity purified
Reconstitution:	Reconstitution with 150 µl of 0.01 M sterile PBS.
	"Note: please spin tube briefly prior to opening it to avoid any losses that might
	occur from lyophilized material adhering to the cap or sides of the tube".
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Storage:	12 months from date of receipt, -20 to -70°C as supplied.
	6 months, -20 to -70°C under sterile conditions after reconstitution.
	1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the
	temperature recommended above.

## **Application Information**

<b>Recommended Dilution:</b>	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	33 kDa




Predicted Reactivity:

Among species analyzed, the sequence of the synthetic peptide used for immunization is 80-99% homologues with the sequence in *Brassica rapa, Brassica napus.* 

For more species homologues information, please contact tech support at <u>tech@phytoab.com</u>.

## Application Example



Recom: 2.5 ng, 15 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 42 kDa.
Electrophoresis: 12% SDS-PAGE
Transfer: blotting to NC (nitrocellulose) membrane for 1 h.
Blocking: 5% skim milk at RT or 4°C for 1 h.
Primary antibody: 1:1000 dilution overnight at 4°C.
Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).
Detection: using chemiluminescence substrate and image were captured with CCD camera.

**Research Use Only**