

# Anti-AT5G40660 antibody

Catalog: PHY2885A

#### **Product Information**

**Description:** Rabbit polyclonal antibody

**Background:** P12 is an F-type ATP Synthase Assembly factor that binds to beta subunits of

mitochondrial ATPase.

Synonyms: P12

Immunogen: KLH-conjugated synthetic peptide (18 aa from Central section) derived from

Arabidopsis thaliana P12 (AT5G40660).

Form: Lyophilized

**Quantity**: 150 μg

Purification: Immunogen affinity purified

Reconstitution: Reconstitution with 150 µl of 0.01 M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability & Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

**Storage:** 12 months from date of receipt, -20 to  $-70^{\circ}$ C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8℃ under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

### **Application Information**

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

**Expected / apparent MW:** 36 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used


for immunization is 80-99% homologues with the sequence in *Vitis vinifera*, *Glycine max*, *Gossypium raimondii*, *Populus trichocarpa*, *Brassica rapa*, *Brassica napus*, *Spinacia oleracea*, *Cucumis sativus*, *Medicago truncatula*, *Hordeum vulgare*, *Triticum aestivum*, *Nicotiana tabacum*, *Solanum lycopersicum*, *Solanum tuberosum*, *Oryza sativa*,



#### Panicum virgatum.

For more species homologues information, please contact tech support at <a href="mailto:tech@phytoab.com">tech@phytoab.com</a>.

## **Application Example**



Recom: 3.75 ng, 22.5 ng and 37.5 ng recombinant protein containing the

peptide for immunization and having a molecular mass of 45 kDa.

Electrophoresis: 12% SDS-PAGE

**Transfer:** blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or 4℃ for 1 h.

**Primary antibody:** 1:1000 dilution overnight at 4℃.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP)

(Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.