

Anti-Zinc-finger homeodomain protein 8 antibody

Catalog: PHY7422S

Product Information

Description: Rabbit polyclonal antibody

Background: HB30 is a member of the zinc finger homeodomain transcriptional factor family.

Synonyms: HB30, ATHB30, HOMEOBOX PROTEIN 30, ZFHD3, ZHD8, ZINC FINGER

HOMEODOMAIN 3, ZINC FINGER HOMEODOMAIN 8

Immunogen: KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from

Arabidopsis thaliana HB30 (AT5G15210).

Form: Lyophilized

Quantity: $150 \mu g$ Purification: Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

Reconstitution: Reconstitution with 150µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70° as supplied.

6 months, -20 to -70 $^{\circ}$ C under sterile conditions after reconstitution.

1 month, 2 to 8℃ under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 29 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used

for immunization is 80-99% homologues with the sequence in

Spinacia oleracea, Brassica napus, Brassica rapa, Vitis vinifera,


Solanum tuberosum, Solanum lycopersicum.

The sequence of the synthetic peptide used for immunization is 80%



(12/15) homologues with the sequence in HB23 (AT5G39760). For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 44 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4℃.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP)

(Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with

CCD camera.