

Anti-Cycloartenol-C-24-methyltransferase, N-terminal antibody

Catalog: PHY3152A

Product Information

Description:	Rabbit polyclonal antibody
Background:	Sterol methyltransferase (SMT), the enzyme from Saccharomyces cerevisiae
	that catalyzes the conversion of sterol acceptor in the presence of AdoMet to
	C-24 methylated sterol and AdoHcy. Arabidopsis contains three SMT, sterol
	methyltransferase 1 (SMT1, AT5G13710), SMT2 (AT1G20330), and SMT3
	(AT1G76090).
Synonyms:	SMT1, CEPHALOPOD, CPH, STEROL METHYLTRANSFERASE 1
Immunogen:	KLH-conjugated synthetic peptide (14 aa from N terminal section) derived from
	Arabidopsis thaliana SMT1 (AT5G13710).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Immunogen affinity purified
Reconstitution:	Reconstitution with 150 µl of 0.01 M sterile PBS.
	"Note: please spin tube briefly prior to opening it to avoid any losses that might
	occur from lyophilized material adhering to the cap or sides of the tube".
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Storage:	12 months from date of receipt, -20 to -70 $^\circ C$ as supplied.
	6 months, -20 to -70 $^\circ\!\!\!\!\!^\circ$ under sterile conditions after reconstitution.
	1 month, 2 to 8 $^\circ C$ under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4 $^\circ\!\!{ m C}$. Upon receipt, store it immediately at the
	temperature recommended above.

Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	38 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used
	for immunization is 80-99% homologues with the sequence in

Research Use Only


PhytoAB Inc.



Brassica rapa, Brassica napus, Glycine max, Vitis vinifera, Cucumis sativus, Gossypium raimondii.

For more species homologues information, please contact tech support at <u>tech@phytoab.com</u>.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 48 kDa.
Electrophoresis: 12% SDS-PAGE
Transfer: blotting to NC (nitrocellulose) membrane for 1 h.
Blocking: 5% skim milk at RT or 4°C for 1 h.
Primary antibody: 1:1000 dilution overnight at 4°C.
Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).
Detection: using chemiluminescence substrate and image were captured with CCD camera.

Research Use Only