

Anti-Photosystem I reaction center subunit II, chloroplastic, C-terminal antibody

Catalog: PHY4491S

Product Information Description: Rabbit polyclonal antibody Background: Os08g0560900 Synonyms: Os08g0560900 Immunogen: KLH-conjugated synthetic peptide (17 aa from C terminal section) derived from Oryza sativa Os08g0560900. Form: Lyophilized Quantity: 150 µg **Purification:** Serum Peptide affinity form antibody available upon request at info@phytoab.com. **Reconstitution:** Reconstitution with 150 µl of sterile water. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube". Stability & Use a manual defrost freezer and avoid repeated freeze-thaw cycles. Storage: 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution. The product is shipped at 4 °C. Upon receipt, store it immediately at the Shipping: temperature recommended above. Application Information **Recommended Dilution:** Western Blot (1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the

Expected / apparent MW:end user.Expected / apparent MW:22 kDaPredicted Reactivity:Among species analyzed, the sequence of the synthetic peptide used
for immunization is 100% homologues with the sequence in
Arabidopsis thaliana, Brassica napus, Brassica rapa, Gossypium
raimondii, Zea mays, Physcomitrium patens, Glycine max, Nicotiana


PhytoAB Inc.

Research Use Only



tabacum, Medicago truncatula, Solanum lycopersicum, Solanum tuberosum, Triticum aestivum, Hordeum vulgare, Populus trichocarpa, Vitis vinifera, Setaria viridis, Sorghum bicolor, and 80-99% homologues with the sequence in *Spinacia oleracea*, *Synechocystis sp.* PCC 6803, *Chlamydomonas reinhardtii*. For more species homologues information, please contact tech support at <u>tech@phytoab.com</u>.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 43 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4° C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.

Research Use Only