

Anti-Magnesium-chelatase subunit Chll-1, chloroplastic, N-terminal antibody

Catalog: PHY3306A

Product Information

Description:	Rabbit polyclonal antibody
Background:	CHLI1 is required for chlorophyll biosynthesis. It was suggested that the redox
	state of CHLI is regulated in vivo by the change of the redox environment in
	the chloroplasts probably via the Trx system.
Synonyms:	CHLI1, CHL11, CH-42, CH42, CHLI-1, CHLORINA 42, LOST1, LOW
	TEMPERATURE WITH OPEN-STOMATA 1
Immunogen:	KLH-conjugated synthetic peptide (15 aa from N terminal section) derived from
	Arabidopsis thaliana CHLI1 (AT4G18480).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Immunogen affinity purified
Reconstitution:	Reconstitution with 150 µl of sterile 1XPBS (PH=7.4).
	"Note: please spin tube briefly prior to opening it to avoid any losses that might
	occur from lyophilized material adhering to the cap or sides of the tube".
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Storage:	12 months from date of receipt, -20 to -70 $^\circ C$ as supplied.
	6 months, -20 to -70 $^\circ$ C under sterile conditions after reconstitution.
	1 month, 2 to 8 $^{\circ}$ C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4 $^\circ\!\!{ m C}$. Upon receipt, store it immediately at the
	temperature recommended above.

Application Information


Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	46 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used
	for immunization is 80-99% homologues with the sequence in <i>Brassica</i>
	napus, Brassica rapa.

PhytoAB Inc.



For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 50 kDa. Electrophoresis: 12% SDS-PAGE Transfer: blotting to NC (nitrocellulose) membrane for 1 h. Blocking: 5% skim milk at RT or 4°C for 1 h. Primary antibody: 1:1000 dilution overnight at 4°C. Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000). Detection: using chemiluminescence substrate and image were captured with CCD camera.